ISSN: 1816-949X

© Medwell Journals, 2017

## Forecasting Analysis Vacant Land Detection

Warnia Nengsih Department of Computer, Politeknik Caltex Riau, Umban Sari Street, Rumbai, 28265 Pekanbaru Riau, Indonesia

**Abstract:** Forecasting analysis is a prediction for possible a condition in the future. The scheme or the knowledge that acquired can be used as a reference for policy or decision making. The reasoning for choice the locations should have good enough knowledge. The difficulty of determine which location's point will be attracted to people in the utilization of the next vacant land also being a foundation in forecasting method and forecasting analysis used. The methods will be used are multiple regression as for values using three variable influential (x) and 1 variable affected (y) to form equation value gradient and constant. Equation value gradient produced is 29.5792 with the constant 1 (0.006459), value constant 2 (0.00657 and value constant 3(0.00847), so that, the prediction of the yielded become a reference to determine locations favorite ones.

Key words: Forecasting analysis, multiple regression, vacant land, constant, variable, Riau

#### INTRODUCTION

The difficulty of determine which location's point will be attracted to people in the utilization of the next vacant land also being a foundation in forecasting method and forecasting analysis used. The forecasting method will be used is multiple regression. Multiple regression is one of predictive models in data mining uses which is predict a value based on value it already is.

The object of this research Pekanbaru, a city which has area  $632.26 \text{ km}^2$  consist of 12 districts with 897.767 population in 2010. This location will be object of research in 58 points. Where representation of this point is a small part of the region. As for variable will be used are 3 independent variables and 1 dependent variable with details  $X_1$  = vacant land uses;  $X_2$  = bulding land uses;  $X_3$  = location point enthused.

With forecasting analysis is expect to detection land's problem can be solved, so that, the schema or the trend obtained can faciltate the public, investor and government to predict location point enthused.

## MATERIALS AND METHODS

## **Theoritical**

**Vacant land:** According to Law of Spatial Planning (UUPR) Constitution of the Republic of Indonesia No. 24/92 about spatial planning as a whole region of human and other things and maintain their survival. And according to the basis of modern Indonesian land law

is Law Number 5 of 1960 (UU 5/1960) also referred to as UUPA (Undang-Undang Pokok Agraria or the Basic Agrarian Law Act) regulate the management land include by the arrangement land uses, tenure of land arrangement, obtaining rhe right land and registration the land (Hasni, 2008).

Based to the FAO in Luthfi (2007), land have so much fuctions of them production function, environment biotic function, officers climate fuction, storage function and spasial connected function.

Multiple regression linear methods: Regression method is one of predictive models. This method usually used to know a schema or trend in a case. And are usually can be use in forecasting analysis. This method's concept use data learning that gained in the exist data. There are several technique in this method, first linear regression and multiple regression (Beasley, 2008).

There is a continuous random variable called the dependent variable, Y and a number of independent variables,  $X_1$ ,  $X_2$ , ...,  $X_n$ . The purpose is to predict the value of the dependent variable (also referred to as the response variable) using a linear function of the independent variables (Beasley, 2008). The values of the independent variables (also referred to as predictor variables, regressors or covariates) are known quantities for purposes of prediction (Zellner, 1962), the model is:

$$Y = a + b_1 X_1 + b_2 X_2 +, ..., + b_n X_n$$
 (1)




Fig. 1: Forecasting analysis diagram

Where:

Y = The dependent variables

A = The constant  $b_1, b_2$  = The gradient

 $X_{1, X2}$  = The independent variebles

The formula equation formed, uses the matric with the reference explanatory variable used.

**Research design:** Forecasting analysis that produced to do some testing for determine the accuracy value and level of errors percentage (Culpepper, 2008). The following diagram forecasting analysis block as seen in Fig. 1.

Figure 1 explain the prediction in general. Start from preprocessing vacant land data then processed it use multiple regression to find predictive value. Next, productive value produced to do some accuracy testing and error rates. As for variable will be used are 3 independent variables and 1 dependent variable with details  $X_1$  = vacant land uses;  $X_2$  = bulding land uses;  $X_3$  = location point enthused. To find equation line preconceived previously, should find the constant and gradient value firstly. If those values have been obtained from equation, so, Y values of that equation can be displayed.

#### RESULTS AND DISCUSSION

As for variable will be used are 3 independent variables and 1 dependent variable with details  $X_1$  = vacant land uses;  $X_2$  = bulding land uses;  $X_3$  = location point enthused which is Y = location's point enthused from 58 records processed data.

Column 1 in Table 1, explain that 3 variable X and 1 variable 1 in used. That variables have been processed use multiple regression method for obtain the equation to determine each predictor value in location point. Column 4 shown the result predictor value and processed result.

Table 1: Data and prediction results										
$X_1$	$X_2$	$X_3$	Y	Prediction results						
1.05	12.600	12.60	1	29.609979830						
1.51	1.580	1.58	2	29.591960340						
0.64	2.000	2.00	3	29.587141180						
0.37	2.300	2.30	4	29.585968730						
7	53.000	53.00	5	29.725362360						
9	66.000	66.00	6	29.763041960						
14	145.000	145.00	7	29.945812680						
7	74.000	74.00	8	29.765362490						
104.28	360.230	360.23	9	30.938876970						
113.34	421.650	421.65	10	31.114384620						
275.75	850.240	850.24	11	32.979723960						
173.69	545.300	545.30	12	31.739698170						
32.9	101.100	101.10	13	29.984264800						
921	2425.000	24.25	14	19.811623600						
779	2857.000	28.57	15	16.094732670						
452	1061.000	10.61	16	25.622364350						
684	3751.000	37.51	17	9.687238855						
3.36	5.320	5.32	18	29.611032960						
35.45	12.840	12.84	19	29.832619920						
16.08	12.890	12.89	20	29.707608080						
16.62	17.700	17.70	21	29.720257770						
3.49	5.100	5.10	22	29.611453560						
0	388.000	388.00	23	30.318248030						
10	215.000	215.00	24	30.053311220						
0	900.000	900.00	25	31.293489300						
0	400.000	400.00	26	30.341105250						
0	209.000	209.00	27	29.977294540						
0	4000.000	4000.00	28	37.198270430 29.727682890						
5 6.75	61.000 44.250	61.00 44.25	29 30	29.707080940						
3	26.000	26.00	31	29.648098400						
1.2	22.8.0	22.80	32	29.630377290						
2	23.000	23.00	33	29.635925290						
2.25	28.750	28.75	34	29.648492410						
13.01	73.990	73.99	35	29.804160870						
5.31	115.690	115.69	36	29.833856890						
3.24	114.760	114.76	37	29.818715730						
24	135.000	135.00	38	29.991353060						
31	123.000	123.00	39	30.013707490						
8.5	59.500	59.50	40	29.747431570						
5.5	62.500	62.50	41	29.733769450						
22.5	96.500	96.50	42	29.908331280						
12	85.000	85.00	43	29.818608980						
1.6	59.400	59.40	44	29.702675320						
1.02	53.980	53.98	45	29.688605370						
1.85	91.100	91.10	46	29.764671170						
2.97	36.030	36.03	47	29.667009460						
1.3	37.700	37.70	48	29.659404210						
1.25	42.800	42.80	49	29.668795590						
1.27	42.730	42.73	50	29.668791430						
269	2090.000	2090.00	51	35.297582320						
286	1083.000	1083.00	52	33.489280550						
302	907.000	907.00	53	33.257382270						
243	801.000	801.00	54	32.674407260						
0	40.010	40.01	55	29.655407780						
0	6.800	6.80	56	29.592150430						
0	6.600	6.60	57	29.591769480						
0	85.600	85.60	58	29.742246160						

Figure 2 shown the process of multiple regression method and do some data reading for split data and role data. And the next step, will be gained all subsets regression (Fig. 3).




Fig. 2: Processed of multiple regression method




Fig. 3: The equation value

Table 2: Output of StatCalc for each atribute

Coefficient	SE	Std. coefficient	Tolerance	t-stat	p-values	Code
-0.005	0.003	-0.258	0.608	-1.529	0.132	
0.0070	0.005	0.263	0.608	1.560	0.124	
29.760	2.484	-	-	11.980	0.000	oje oje oje oje

Table 2 shown the accuracy testing for each atribute in used from linier regression produced. Next, prediction testing for each location in aplication above with testing variable uses. Data training aplication to get prediction result of the equation had been formed. Where the equation:

$$Y = 29.5792 + 0.006459 X_1 - 0.00657 X_2 + 0.00847 X_3$$

with the gradient produced is 29.5792, constant value 1 (0.006459, constant value 2 (0.00657 and constant value 3 (0.00847). The prediction value in produced will be the trend or reference to determine the location favorite ones (Fig. 4).

Fig. 4: Data training aplication

#### CONCLUSION

Forecasting analysis is a prediction for possible a condition in the future. The scheme or the knowledge that acquired can be used as a reference for policy or decision making. The reasoning for choice the locations should have good enough knowledge.

# REFERENCES

Beasley, T.M., 2008. Seemingly unrelated regression SUR models as a solution to path analytic models with correlated errors. Multiple Linear Regression Viewpoints, 34: 1-7.

Culpepper, S.A., 2008. Conducting external profile analysis with multiple regression. Pract. Assess. Res. Eval., 13: 1-8.

Hasni, 2008. [Law on Spatial Planning and Land Use Administration: In the Context of the BAL-UUPR-UUPLH]. Penerbit Rajawali Pers, Jakarta, Indonesia, (In Indonesian).

Luthfi, R., 2007. [Land Resource Inventory Method]. Penerbit Andi, Yogyakarta, Indonesia, (In Indonesian).

Zellner, A., 1962. An efficient method of estimmating seemingly unrelated regressions and tests for aggregation bias. J. Am. Statist. Assoc., 57: 348-368.